

Induced Violation of Weak Equivalence Principle in the Brans-Dicke Theory

Yasunori Fujii*

Nihon Fukushi University

Okuda, Chita-gun, Aichi, 470-32 Japan

and

Institute for Cosmic Ray Research

University of Tokyo

Tanashi, Tokyo, 188 Japan

Abstract

A quantum correction to the Brans-Dicke theory due to interactions among matter fields is calculated, resulting in violation of WEP, hence giving a constraint on the parameter ω far more stringent than accepted so far. The tentative estimate gives the lower bounds $\gtrsim 10^6$ and $\gtrsim 10^8$ for the assumed force-range $\gtrsim 1\text{m}$ and $\gtrsim 1\text{AU}$, respectively.

*E-mail: ysfujii@tansei.cc.u-tokyo.ac.jp

Many aspects of the Brans-Dicke (BD) theory [1], as a classical theory, have been the focus of extensive studies. Little attention seems to have been paid, however, to quantum effects due to the interaction among matter fields. This might be serious, because it results in violation of Weak Equivalence Principle (WEP)[†] to the extent that the bound on the parameter ω will be made much more stringent than accepted so far, as will be demonstrated on the basis of a simple model.

Assume the Lagrangian:

$$\mathcal{L} = \sqrt{-g} \left[\frac{1}{8\omega} \phi^2 R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi + L_{\text{matter}} \right], \quad (1)$$

where our ϕ , related to the original symbol by $\varphi_{BD} = \phi^2/8\omega$, fits the conventional notation in the standard quantum field theory. The solar system experiments have given a widely accepted lower bound $\omega \gtrsim 500$.

The nonminimal couplings of the scalar field essentially of the same nature are known to be rather common in many of the theoretical models of unification, thus making this classic model particularly relevant.

The assumed masslessness of the scalar field is, however, vulnerable against quantum effects; there is no principle that prevents a scalar field from acquiring a nonzero mass. It is even likely that the acquired mass-squared is $\mu^2 \sim Gm^4$, giving a force-range $\lambda = \mu^{-1}$ which would be of the order of macroscopic distances if m is of the order of hadronic masses. This observation was one of the motivations to propose a non-Newtonian force, or the fifth force [2]. The constraint $\omega \gtrsim 500$ would not apply if $\lambda \ll 1$ AU.

Another important requirement in the original BD theory is the assumed absence of the *direct* coupling of ϕ to the matter fields at the level of the Lagrangian, though the coupling arises *indirectly*. In fact this assumption results in the geodesic equation for a matter particle, hence ensuring WEP. The purpose of this note is to show that this result is also fragile against quantum effects.[‡]

For the sake of illustration we consider the matter system of nucleons and the electromagnetic field. The analysis is focused upon electromagnetic interaction of protons. In the original theory of (1), however, the ϕ -matter coupling occurs only in the field equation. This makes it inconvenient to apply the conventional

[†]WEP, sometimes called composition-independence, implies that any object at a given space-time point falls with a common acceleration independent of its mass and composition. See, for example, C. Will, Phys. Rep. **113**, 346 (1984).

[‡]Violation of WEP does not necessarily imply the same of Equivalence Principle (EP) in the more fundamental and theoretical sense; tangential spacetime at each point of curved spacetime should be Minkowskian.

technique of QED. We then apply a conformal transformation [3][§]:

$$g_{\mu\nu} \rightarrow g_{*\mu\nu} = \frac{2\pi G}{\omega} \phi^2 g_{\mu\nu}. \quad (2)$$

In the new starred conformal frame the Lagrangian (1) is cast into the form

$$\mathcal{L} = \sqrt{-g_*} \left[\frac{1}{16\pi G} R_* - \frac{1}{2} g_*^{\mu\nu} \partial_\mu \sigma \partial_\nu \sigma + L_{*\text{matter}} \right], \quad (3)$$

where σ is a redefined canonical scalar field related to ϕ by

$$\phi = \sqrt{\frac{\omega}{2\pi G}} e^{\beta\sigma}, \quad \text{with} \quad \beta = \sqrt{\frac{4\pi G}{3 + 2\omega}}.$$

For the matter Lagrangian with the proton field ψ and the electromagnetic field A_μ , the transformed Lagrangian differs in form only in the mass term:

$$L_{*\text{mass}} = -m e^{-\beta\sigma} \bar{\psi}_* \psi_*, \quad (4)$$

where $\psi_* = (2\pi G/\omega)^{-3/4} \phi^{-3/2} \psi$, while A_μ remains unchanged. The exponential factor in (4) shows the presence of the direct matter coupling in the new conformal frame. In accordance with this, the geodesic equation is modified but in such a universal manner that WEP remains valid.[¶] In the following we suppress the symbol $*$ to simplify the notation.

We focus upon the linear term

$$L_\sigma = \beta m \bar{\psi} \psi \sigma. \quad (5)$$

This is the coupling to the trace of the matter energy-momentum tensor $T = -m \bar{\psi} \psi$, with no contribution from A_μ . This interaction, corresponding to a vertex in Feynman diagrams, may be represented by a “mass insertion” to a proton line in the limit of a vanishing momentum transferred to σ .

We now calculate the one-loop correction to this vertex. To the lowest-order approximation with respect to the fine-structure constant $\alpha = e^2/4\pi \approx 1/137$, we consider two types of diagrams: (a) mass insertion in the *internal* proton line of the proton self-energy diagram (se); (b) mass insertion to the *external* proton lines attached to (se).

The contribution from the diagram (se) is represented by

$$\Sigma(m, i\not{p}) = A(m) + (m + i\not{p}) B(m) + (m + i\not{p})^2 \Sigma_f(m, i\not{p}),$$

[§]The final results remain the same in both conformal frames before and after the transformation in our calculation to the lowest order with respect to G , though this is not the case in general. The equivalence can be established by introducing σ in the *original* conformal frame; $\phi = \sqrt{\omega/2\pi G} + (1/\sqrt{1+3/2\omega})\sigma$. Diagonalization corresponding to (2) is also useful. More details will be discussed elsewhere.

[¶]The right-hand side of the geodesic equation is now $-\beta(\partial_\nu \sigma)(g_*^{\mu\nu} + u_*^\mu u_*^\nu)$, being independent of properties specific to individual objects.

where $A = \delta m$ is the self-energy as given by a divergent integral [4]

$$A = m \frac{\alpha}{2\pi} \left[-i \frac{1}{\pi^2} \int dx (1+x) \int d^4k \frac{1}{(k^2 + x^2 m^2)^2} - \frac{1}{4} \right], \quad (6)$$

while B represents a wave-function renormalization.

The correction results in replacing m in (5) by $\mathcal{M}$ in which the portion proportional to α is given by $\mathcal{M}_1 = \mathcal{M}_a + \mathcal{M}_b$, where

$$\mathcal{M}_a = \left[\left(m \frac{\partial}{\partial m} \right) \Sigma(m, i\not{p}) \right]_{i\not{p}=-m}, \quad \text{and} \quad \mathcal{M}_b = -mB. \quad (7)$$

In $\mathcal{M}_a$, $(\partial/\partial m)$ adds another proton line, while m provides the mass at the vertex. In the calculation of $\mathcal{M}_b$, on the other hand, we have included $-\delta m$ in the self-energy part. A cancellation occurs among the terms of B , hence leaving

$$\mathcal{M}_1 = m \frac{dA}{dm} = \delta m - \frac{3\alpha}{2\pi} m, \quad (8)$$

where (6) has been used.

The first term δm is obviously absorbed into the zeroth order term m , thus replacing the bare mass by the observed renormalized mass $m + \delta m$. On the other hand, the second term $-(3\alpha/2\pi)m$ represents an extra contribution not to be included in the trace. The fact that this term is not part of mass can be seen by examining T_{00} which is the source of the gravitational field for a *static renormalized* proton:

$$\begin{aligned} T_{00} &\approx \frac{1}{2} \bar{\psi} (\gamma_0 \vec{\partial}_0 - \overleftarrow{\partial}_0 \gamma_0) \psi \\ &= -\frac{1}{2} \bar{\psi} (\vec{\partial} - \overleftarrow{\partial}) \psi \\ &= \bar{\psi} [(m + \delta m) - \beta m \sigma + \dots] \psi, \end{aligned}$$

where $\dots$ means other interaction terms. Notice that due to the Ward identity, no renormalization is present for $T_{\mu\nu}$ as long as we stay out of quantum effects of gravity in 4 dimensions.^{||} We have retained the σ -term because it might behave like a constant in a region where the field $\sigma(\vec{x})$ is nearly constant. This term, however, is negligible of the order of $\sqrt{G}$ being entirely different from the above extra term $-(3\alpha/2\pi)m$. In this way we find that *the extra term gives a difference between what the tensor gravitational field feels and what the scalar field feels, hence the violation of WEP*. We also observe that the term depends explicitly on α , one of the “internal parameters,” beyond the extent to which it is absorbed into the mass.

^{||}I thank K. Fujikawa for the related comments.

We point out that the extra term is *finite* in the same context as in the “trace anomaly”.** WEP which has been designed to be valid classically is induced to be violated following the manner of “quantum anomalies.” This finiteness allows us to make a less ambiguous prediction than the force-range of σ . It is interesting to notice that the finiteness can also be understood by using an explicit expression $\delta m = (3\alpha/2\pi)m[\ln(\Lambda/m) + 1/4]$, where Λ is a cutoff. The anomalous term arises simply because a naive dimensional analysis fails due to the presence of the cutoff.

We now discuss phenomenology. By assuming a nonzero mass $\mu = \lambda^{-1}$ of σ , we have a fifth-force-type potential:

$$V_{ij} = -G \frac{M_i M_j}{r} \left(1 + \frac{q_i q_j}{3 + 2\omega} e^{-r/\lambda} \right), \quad (9)$$

as will be read off from (5).

According to the present model, the neutral neutron is free from anomaly; $q_n = 1$, whereas (8) for the proton is translated into $q_p = 1 - (3\alpha/2\pi)$.

Consider a nucleus i with the mass number A_i and the atomic number Z_i . Again as an illustration, we choose a simplified model in which a nucleus is made of nucleons and fields mediating nuclear forces but assumed to *observe* WEP. We also ignore electrons. Then q_i for the atom/nucleus is given by

$$q_i = 1 - \frac{3\alpha}{2\pi} \frac{Z_i m}{M_i} \approx 1 - \frac{3\alpha}{2\pi} \frac{Z_i}{A_i},$$

where M_i is the mass of the nucleus and the proton mass m replaced by $\approx M_i/A_i$.

The fractional difference of the acceleration between the two nuclei i and j toward a source S is then

$$\frac{\delta a_{ij}}{a_{ij}} = \frac{1}{3 + 2\omega} (q_i - q_j) q_S \mathcal{F},$$

where q_S of the source may be replaced by the composition-independent component 1, while $\mathcal{F}(R, \lambda)$ takes care of the finite force-range; $\mathcal{F} \rightarrow 1$ for $\lambda \gg R$.

Suppose first that $\lambda \gtrsim 1\text{AU}$. Then the most stringent constraint comes from the null experiments by Roll, Krotkov and Dicke, and by Braginski and Panov [5]. These result impose the condition $|\delta a_{ij}/a_{ij}| \lesssim 10^{-12}$. This translates into

$$\left| \frac{1}{3 + 2\omega} \frac{3\alpha}{2\pi} \delta \left(\frac{Z}{A} \right) \right| \lesssim 10^{-12}.$$

We find $\delta(Z/A) \approx 0.08$ for Al and Pt/Au, hence giving $\omega \gtrsim 1.5 \times 10^8$, far larger than $\omega \gtrsim 500$, as obtained from the solar-system experiments.

**Trace anomaly itself is consistent with EP at the fundamental level. It can be relevant to the experimental WEP violation only if σ is sufficiently long-ranged.

If $\lambda \ll 1\text{AU}$, the free-fall experiments on the Earth [6] then give the lower bounds of ω ; 460, 5.2×10^3 and 2.5×10^4 for the assumed values of λ ; 100km, 10^3km and 10^4km , respectively. They still tend to be more stringent than 500.

If λ is as small as 1m or even less, we resort to the composition-dependent experiments using terrestrial sources. Some of the recent measurements [7] reached the accuracy $\sim 10^{-11}$ for the upper bound of the fractional acceleration difference between Cu and Be, giving $\omega \gtrsim 2.2 \times 10^6$. This is true for $\lambda \gtrsim 1\text{ m}$, thus surpassing the limits set by the free-fall experiments.

The amount of composition-dependence as represented by $3\alpha/2\pi$ is quite large as compared with what has been expected from the baryon-number coupling. Combining this with extremely accurate tests of WEP is the reason why we “improved” the bound of ω by many orders of magnitude. The new constraint will affect many analyses attempted so far on the BD theory. On the other hand, very large ω may be avoided if λ is sufficiently small, perhaps shorter than 1m.

Our analysis was based on a simplified model in which composition-dependence comes only from the anomaly in protons. For this reason the results are tentative, mainly to demonstrate the importance of the issue. Obviously other fields should be taken into account. Even the electromagnetic interaction of the neutron through the anomalous magnetic moment may not be ignored. It might be better to apply the same technique to the quarks with the gauge fields at a more fundamental level. With such elaborations, however, the effect of roughly the same size still seems unavoidable unless some cancellation mechanism is discovered to work.

A possible way to avoid unnaturally large values of ω is to invoke a suppression mechanism as advocated by Damour and Nordtvedt [8]. Even with the suggested modification of the nonminimal coupling, however, the anomalous effect should be still present to be included in the analysis. In this connection we also point out that the argument depends crucially on the assumption of a *standard* mass term in the original conformal frame in which the Lagrangian is given by (1). Unification models suggest the ϕ -dependence of the mass terms. An example was studied in which the “mass” tends asymptotically to a constant value as the Universe expands [9].^{††} Further study of various versions of the scalar-tensor theory in conjunction with the experimental efforts will benefit the development toward unification.

Finally we add that the mass insertion to the photon self-energy part yields the anomalous coupling

$$L_{\sigma\gamma} = -\frac{2\alpha}{3\pi}\beta\frac{1}{4}F_{\mu\nu}F^{\mu\nu}\sigma,$$

^{††}According to this model, the strength of the σ coupling is suppressed by the age of the Universe.

which is also finite. This is interesting because it is not a trace coupling, certainly violating WEP. Unfortunately, the effect through the nuclear Coulomb energy is numerically negligible compared with that of the proton anomaly.

I thank Y.M. Cho for discussions at the early stage of the work. I also acknowledge useful discussions with K. Kuroda and R. Newman.

References

- [1] C. Brans and R.H. Dicke, Phys. Rev. **124**, 925 (1961).
- [2] Y. Fujii, Nature Phys. Sci, **234**, 5 (1971); Phys. Rev. **D9**, 874 (1974).
See, for the latest review, E. Fischbach and C. Talmadge, Nature, **356**, 207 (1992).
- [3] R.H. Dicke, Phys. Rev. **125**, 2163 (1962).
- [4] See, for example, J.M. Jauch and F. Rohrlich, *The Theory of Photons and Electrons*, Addison-Wesley, 1954; C. Itzykson and J.-B. Zuber, *Quantum Field Theory*, McGraw-Hill, 1985.
- [5] P.G. Roll, R. Krotkov and R.H. Dicke, Ann. Phys. (N.Y.) **26**, (1964) 442; V.B. Braginski and V.I. Panov, Sov. Phys. JETP **34**, (1972) 463.
- [6] T.M. Niebauer, M.P. McHugh and J.E. Faller, Phys. Rev. Lett. **59**, 609 (1987); K. Kuroda and N. Mio, Phys. Rev. Lett. **62**, 1941 (1989). The results in the text were obtained from the second reference.
- [7] E.G. Adelberger et al., Phys. Rev. **D42**, 3267 (1990).
- [8] T. Damour and K. Nordtvedt, Phys. Rev. Lett. **70**, 2217 (1993); See also, T. Damour and A.M. Polyakov, The String Dilaton and a Least Coupling Principle, preprint.
- [9] Y. Fujii and T. Nishioka, Phys. Rev. **D42**, 361 (1990).